## **Pyrethroid Benefits & Science**

Presented To Central Valley Regional Water Quality Control Board

Ву

Tess Dunham, Somach Simmons & Dunn Michael Dobbs, Bayer Crop Science

On Behalf of The Pyrethroid Working Group (PWG)
August 18, 2016

(c) Pyrethroid Working Group

## **About Pyrethroids**

- PWG is an industry group comprised of pyrethroid manufacturers dedicated to research and stewardship
- PWG has extensively researched
  - From data summaries to modeling to lab to field monitoring studies
  - Have worked to address questions from the regulatory and broader community
- Have been in use for over 35 years
- Most widely used urban insecticides
- Have a favorable human health profile

#### **Benefits**

- **Protects Public Health** 
  - Helps Control Pests that Transmit Serious Illnesses
    - Zika Virus
    - West Nile Virus
    - Malaria
- **Helps Control Pests** 
  - Parasitic: Lice, Bedbugs
  - Nuisance: Cockroaches, ants, spiders
  - Property Damage: Termites
- Food Crops
  - Control of wide range of pests





\*ADAM





#### **Examples of PWG Research**

#### **PWG Pathway ID Study**

- Validated the
   effectiveness of
   revised label
   restrictions and DPR
   regulations to
   substantially mitigate
   pyrethroid runoff
  - 40-fold reduction

#### **Biomonitoring Studies**



- Pyrethroids generally not the primary stressor impacting the benthic community
- Over the course of 10 years, Pleasant Grove Creek showing a declining trend of pyrethroids concentrations
- Hall, Anderson, and Killen, Arch Environ Contam Toxicol
   (2016) 70:332–340 (c) Pyrethroid Working Group

# Questions raised by Water Board Members February 2016


- Food Chain Interactions
- Bioaccumulation
- Bioagnification
- Sublethal effects
- Sediment ingestion

#### **Food Chain Interactions**


- Most fish are opportunistic feeders
  - they feed on whatever species are most abundant
- While a few crustacean species (e.g. lab-reared *Hyalella azteca*) are extremely sensitive to pyrethroids, other crustaceans as well as insects, worms, and mollusks are much less sensitive
- The likelihood of fish survival, growth, and reproduction being affected by small changes in overall invertebrate abundance is very low

#### Evidence from mesocosm studies

 More than 50 mesocosm studies with pyrethroids show that effects of realistic exposure levels on community structure and productivity are minimal.







## Bioaccumulation/Biomagnification

- Despite being highly adsorptive, pyrethroids do not bioaccumulate significantly in aquatic organisms
- Unlike some legacy highly adsorptive chemicals, pyrethroids are metabolized and depurated rapidly therefore bioconcentration is <1000</li>
  - the usual criterion for bioaccumulative compounds.
- In addition, when modeled using EPA's standard tools,
   BCFs are generally predicted to be less than 1000
- Pyrethroids do not bioaccumulate because they are metabolized and excreted by organisms
- Because pyrethroids do not accumulate in aquatic organisms, they do not biomagnify through the food chain

## Sublethal effects of pyrethroids are accounted for in standard risk assessments

- A wide variety of sublethal effects on aquatic organisms have been studied with pyrethroids
  - From biochemical to behavioral
- Most sublethal effects have not been quantitatively linked to survival, growth, and reproduction
- Concentrations reported to cause sublethal effects are similar to standard acute and chronic toxicity endpoints
  - Concentrations are significantly higher than the proposed trigger values.

## Sediment ingestion is not a significant route of exposure of aquatic organisms to pyrethroids

- Equilibrium partitioning (EqP) theory: toxicity of hydrophobic chemicals is caused by freely dissolved chemical in the pore water (ie. Bioavailability)
- Applicability of EqP to pyrethroids has been well established
- These observations imply that pyrethroids sorbed to sediment are not a significant source of exposure

## Summary


- Pyrethroids **do not** bioaccumulate significantly or biomagnify through the food chain
- Sublethal effects are accounted for in standard risk assessments
- Sediment ingestion is not a significant route of exposure for aquatic organisms.
- Bioavailability is a critical consideration in establishing and interpreting WQC for pyrethroids.

### **Pyrethroids Resource Website**

Developing an online resource for information seekers interested in learning more

month (September 2016)

information beneficial to your understanding of pyrethroids.



# Input on the Strawman Regulatory Approach

- Appreciate efforts to engage stakeholders
- Appreciate alternative approaches for implementation of water quality criteria
- Prefer use of 5<sup>th</sup> percentile triggers over 1<sup>st</sup> percentile values
- Support consideration of bioavailability
- Support phased approach

## **QUESTIONS?**